

The Angular-momentum aspect of *Shift* in FQHE

Dipti Banerjee
Department of Physics
Rishi Bankim Chandra College
Naihati,24-Parganas(N)
Pin-743 165, West Bengal
INDIA

Abstract

The angular momentum of the Hall particles acquiring Berry phases from quantization, is the very source of unifying the physics of IQHE and FQHE. The origin of *Shift* quantum number lies in the change of angular momentum of the Hall particles from higher to lower Landau level. Here the non-vanishing effect of *Shift* in 2D has been evaluated in the framework of Z_p spin system and in view of these, Ising model and XY model are studied.

In a magnetic field B , when a particle is transported over a closed contour, a non-integral phase is developed known as Dirac phase that is determined by the flux through this contour[1].

$$\phi_D = \frac{e}{c\hbar} \int A^\alpha dr_\alpha = \frac{e}{c\hbar} \int F^{\alpha\beta} dS_{\alpha\beta} = \frac{e}{c\hbar} \int B dS = \frac{e}{c\hbar} \Phi \quad (1)$$

This phase of Dirac is similar to the Berry phase [2] when the eigenvectors of a particle are described by the parameter dependent Hamiltonian. The nontrivial connection A_μ under the parallel transport of the eigenvectors acquire the phase.

$$\phi_B = \oint A_\mu d\mu = \int F_{\mu\mu} ds \quad (2)$$

The curvature $F_{\mu\mu}$ associated with Berry connection A_μ turns out to be nonzero in the adiabatic approximation and occurs for loops in the μ space in the manner as the magnetic field B does. This gauge field is the origin of the Chern-Simons term in both 2D and 3D cases. The effect of the Chern-Simons term is to associate with each particle a magnetic flux e/μ . It may be viewed as if a certain number of flux quanta is attached to each electron leading to a gauge theoretic extension. This is the picture of composite particles in condensed matter Physics. This CS term itself generates a phase factor, in addition to the one generated by the Aharonov-Bhm factors associated with $\int d^3x \vec{j} \cdot \vec{A}$ [3]. There is a significant difference between the statistics of ordinary charge/flux tube composites, and that of objects where the charge-flux relation is determined dynamically by the equation of motion of the gauge field. In fact it is the induced charge that is proportional to the induced flux.

$$\int q_{ind}(\Phi) d\Phi$$

As a result the amount of charge present is less than the final amount and the resulting angular momentum is less than before.

This above analysis is reflected in the works of Bandyopadhyay[4] where it has been shown that the charged particle in the 3D anisotropic space (in an intense magnetic field) has the conserved angular momentum

$$\vec{J} = \vec{r} \times \vec{p} - \mu \vec{r} \quad (3)$$

where μ is associated with the induced effect of flux in the angular momentum.

In a Landau problem, when 2D electron gas is taken to reside on the surface of a 3D sphere of large radius R in a radial (monopole) strong magnetic field, particles achieve quantization acquiring the Berry phase through chiral symmetry breaking[6]. The situation arises when the system $\vec{H}_o$ interacts with certain internal Hamiltonian h_{int} to have $\vec{H} = \vec{H}_o + h_{int}$. The response of the internal system corresponding to the change of the external canonical system becomes a part of the action as to associate the change of fields by[7]

$$S_{eff} = S_o + \hbar \Gamma(C) \quad (4)$$

The geometrical phase $\Gamma(C)$, is analogous to the magnetic flux associated with the induced vector potential $A = i < 0|d|0 >$ which emerges from Dirac pole. This eventually causes the deformation of the symplectic structure and the canonical variables originally defined are no longer canonical.

It has been pointed out by Bandyopadhyay [8] that the Berry phase may be associated with the deformation of the symplectic structure which can be linked up with the various quantization procedure. Indeed the relativistic generalization of the quantization procedure can be formulated in terms of gauge theoretic extension of the relativistic quantum particle. The inclusion of constraint of area preserving diffeomorphism in the Landau problem of $2D$ electron gas may be considered as the magnetic translations leading to the quantization of the system. This suggests that there is a dual relationship between the deformation parameter q of the symmetry group (algebra) with the Berry phase parameter μ where the phase is given by $\exp 2\pi i\mu$. Bandyopadhyay pointed out that the deformation through the anisotropy in space-time can be realized in any dimension, with the connections between the Chern-Simons term in the $(2+1)$ dimension, the chiral anomaly in $(3+1)$ and the conformal field theory in $1+1$ -dimension.

In the light of Haldane [9], we first consider the statistical interaction in $3+1$ dimension. Thus, the $2D$ Hall surface is a boundary of a $3D$ sphere having radius R , in a radial (monopole) magnetic field $B = \frac{\hbar S}{eR^2}$. This $2S = N_\phi$ is an integer which defines the total number of magnetic flux through the surface. For parent state and the hierarchical state, the respective fluxes are

$$S = 1/2m(N-1)$$

$$S = 1/2m(N-1) \pm 1/2\left(\frac{N}{p} + 1\right)$$

These show that the filling factor satisfies a slightly deviated relation as

$$2S = N_\phi = \nu^{-1}N - \mathcal{S} \quad (5)$$

Physicists identified this $\mathcal{S}$ as *Shift*, a topological quantum number developed due to the coupling between the orbital spin and curvature of the space. This orbital spin is different from the spin of electrons and is associated with orbital angular momentum in cyclotron motion. In thermodynamic limit, this *Shift* becomes insignificant leading the filling factor in each Landau level by $\nu = \frac{N}{2S} = \frac{N}{N_\phi}$.

In this anisotropic space, the chiral symmetry breaking originates from the curvature $F_{\mu\nu}$ arises from induced flux

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu] \quad (6)$$

The vector potential $A_\mu \in SL(2C)$ is responsible for the deformation which can be linked up with the chiral anomaly. This non-dynamical gauge A_μ is associated with the flux which in $2+1$ dimension is the very cause of Chern-Simon term appeared in the Lagrangian.

$$L_{CS} = \frac{\mu}{2} \epsilon^{\mu\nu\lambda} A_\mu \partial_\nu A_\lambda \quad (7)$$

This licenses a conservation of topological current J_μ which include a topological invariant term in the $(2+1)$ dimension

$$H = \frac{\theta}{2\pi} \int d^3x A_\mu J^\mu \quad (8)$$

in the action. In fact it is the Hopf invariant describing basic maps of S^3 to S^2 . If ρ denotes a four dimensional index then we find [5]

$$\partial_\rho \epsilon^{\rho\mu\nu\lambda} A_\mu F_{\mu\nu} = \frac{1}{2} \epsilon^{\rho\mu\nu\lambda} F_{\rho\mu} F_{\nu\lambda} \quad (9)$$

which connects the Hopf invariant with chiral anomaly. This Hopf term plays a role somewhat similar to the role played by the Wess-Zumino interaction in connection with $3+1$ dim. Skyrmin term.

We have pointed out earlier[10] that the quasi-particles in the hierarchies are formed when additional flux is attached with quantized particles, which in the effective theory is visualized through the additional CS terms in the Lagrangian on $(3+1)$ dimension. Thus quantization is the very cause of anisotropy on the Hall surface along with the internal anisotropy of the particles. Apart from the gauge extension A_μ induced by the external strong magnetic field strength $F_{\mu\nu}$, each particle have its own internal gauge $C_\mu \in SL(2C)$ which is visualized through the field strength $\tilde{F}_{\mu\nu}$.

$$\tilde{F}_{\mu\nu} = \partial_\nu C_\mu - \partial_\mu C_\nu + [C_\mu, C_\nu] \quad (10)$$

These two gauges A_μ and C_μ act as fibres at each point of base space occupied by Hall particles. Hence the quantization of particles in Hall hierarchies involve gauges which can be visualized through their change of coordinates from x_μ to $x_\mu + i(A_\mu + C_\mu)$.

On the surface of 3D sphere the chiral symmetry breaking of composite fermion in Hall fluid is associated with internal and external gauge fields $F_{\mu\nu}$ and $\tilde{F}_{\mu\nu}$ and their physics can be described by the topological Lagrangian [10]

$$\mathcal{L} = -\frac{\theta}{16\pi^2} Tr^* F_{\mu\nu} F_{\mu\nu} - \frac{\theta'}{16\pi^2} Tr^* F_{\mu\nu} \tilde{F}_{\mu\nu} - \frac{\theta''}{16\pi^2} Tr^* \tilde{F}_{\mu\nu} \tilde{F}_{\mu\nu} \quad (11)$$

Here every term corresponds to a total divergence of a topological quantity, known as Chern-Simons secondary characteristics class defined by

$$\Omega^\mu_e = -\frac{1}{16\pi^2} \epsilon_{\mu\nu\alpha\beta} Tr [A_\nu F_{\alpha\beta} - 2/3 (A_\nu A_\alpha A_\beta)] \quad (12)$$

$$\tilde{\Omega}^\mu = -\frac{1}{16\pi^2} \epsilon_{\mu\nu\alpha\beta} Tr [C_\nu F_{\alpha\beta} - 2/3 (C_\nu A_\alpha A_\beta)] \quad (13)$$

$$\Omega^\mu_i = -\frac{1}{16\pi^2} \epsilon_{\mu\nu\alpha\beta} Tr [C_\nu \tilde{F}_{\alpha\beta} - 2/3 (C_\nu C_\alpha C_\beta)] \quad (14)$$

Assuming a particular choice of coupling $\theta = \theta' = \theta''$ in the Lagrangian the topological part of the action in (3+1) dimension become

$$W_\theta = 2(\mu_e + \mu_i + \tilde{\mu})\theta \quad (15)$$

It gives rise the topological phase of Berry on the parallel transport over a closed path of a Hall hierarchy state.

$$\phi_B = \pi W_\theta = 2\pi\tilde{\mu}_{eff}\theta = 2\pi(\mu_e + \mu_i + \tilde{\mu})\theta \quad (16)$$

Here the first term μ_e is associated with Berry Phase (BP) factor due to the anisotropy in the external space. The second term gives rise to the inherent BP factor μ_i associated with the chiral anomaly of a free electron for its internal anisotropy and the third one $\tilde{\mu}$ effectively relates the coupling of the external field with the internal one. It forces the Hall particles in the presence of perpendicular magnetic field to move around the quantized circular orbit. The angular momentum of the rotating Hall particles varies with the radius of the orbit. Hence BP varies directly with angular momentum. With these views, we in our previous works are able to unify the physics of IQHE, FQHE [5] through this angular momentum factor μ of the Hall particles acquiring Berry Phase by $2\pi\mu i$ having $\mu = 0, \pm 1/2, \pm 1, \pm 3/2, \dots$. Further their hierarchies are studied [10] through the angular momentum aspect of Berry's topological phase. Hence the topological phase of the quantized Hall quasi-particles is

$$\phi_B = 2\pi(\mu_{eff} + \mathcal{S})\theta \quad (17)$$

representing the solid angle subtended by their gauges (acting as vortex) with the axis of quantization about which the composite-particles precesses. This is a 3D manifestation of the topological phase. On 2D the Berry Phase is manifested through angles. The *Shift* extends these angles by introducing some deformations.

It has been pointed out earlier [10] that the μ_{eff} appeared in topological phase (eqn.17) actually visualizes the filling factor through the relation $\nu = \frac{n'}{2\mu_{eff}}$, where n' denotes the n' th Landau level. In fact this μ_{eff} satisfies the Dirac quantization condition

$$e'\mu_{eff} = \pm \frac{n'}{2} \quad (18)$$

where $n' = 1, 2, 3, \dots$ denotes the hierarchy levels. Each quasiparticle in the n' th Landau level having charge e' behaves as a composite fermion. It will behave as fermion only when

$$\tilde{e}(\mu_{eff} \mp \frac{n' \pm 1}{2}) = \pm \frac{1}{2} \quad (19)$$

Here the $+$ ($-$) sign indicates the orientation of the vortex line. The $(n' \pm 1)/2$ is the magnetic strength μ of the additional quanta which is associated with the composite fermion in the higher Landau level to change it fermion in ground level. For $\mu = \pm 1/2, \pm 3/2, \dots$ the quanta behaves like fermion and $\mu = \pm 1, \pm 2, \dots$ it

shows bosonic behavior. Hence the behavior of composite particle will be fermionic provided it follows

$$\tilde{e}\tilde{\mu} = \pm \frac{1}{2} \quad (20)$$

having charge $\tilde{e}$ and flux $\tilde{\mu}$. These composite particles satisfying eqn.(20) not always an elementary fermion possessing unit electronic charge ($e = 1$) and magnetic charge ($\mu = \pm 1/2$). *Shift* is visualizing this change of magnetic charge by

$$2\tilde{\mu} = \mathcal{S} = 2\mu_{eff} \mp (n' \pm 1) \quad (21)$$

In other words,

$$\mathcal{S} = J_{CF} \mp (n' \pm 1) \quad (22)$$

where the angular momentum of the composite fermion $J_{CF}=2\mu_{eff}$ is associated with the attached induced flux μ_e in addition with internal flux μ_i . It may be noted that the angular momentum of a quantized Hall particle is $J = \mu_i$ and following Haldane [9] the Landau filling factor $\nu = 1/m$ of parent state in the Hall fluid is the relative angular momentum $m = J_{ij} = J_i + J_j$ of two Hall particles [5].

With these ideas the filling factor of the composite particle is

$$\nu_{CF} = \frac{n'}{2\mu_{eff}} = \frac{n'}{J_{CF}}$$

and leads us to realize $\mathcal{S}$ by the fermionic filling factor whose relationship with the composite fermionic behavior of the Hall particles becomes.

$$\nu_F^{-1} = n'\nu_{CF}^{-1} \mp (n' \pm 1) \quad (23)$$

In 2D Hall surface, Wen [11] pointed out vanishing effect of *Shift*. It seems to us that the effect of *Shift* will not vanish completely on 2D, since anisotropy incorporates some deformation of symplectic structure on Hall surface visualized through curvature of space-time. This physics can be recast through the representation of Hall particles as Z_p spin system. Indeed, the Z_p spin model can be well described by a system of scalar particles, each having an orbital angular momentum $J = 1/p$, can form only one of the p discrete angles $\theta_m = 2\pi m/p$ with some fixed direction in the space of internal degrees of freedom. This gives rise to the behavior of such particles having fractional statistics [12]. An equivalent relationship between the anyon statistical formulation of the hierarchical states depicted in the continued fraction scheme and Jain classification scheme based on the composite fermion model has been derived [13] in the background of Z_p spin system.

The statistics of the particles θ in this Z_p spin system is related to the angular momentum by J [3].

$$\exp^{2\pi i J} = \exp^{i\theta} \quad (24)$$

Here the statistical parameter is represented by

$$\theta = \pi[1 - \frac{1}{n}] \quad (25)$$

where $n = \infty$ corresponds to a fermion and $n = 1$ a boson leading to their respective statistics $\theta = \pi$ and $\theta = 0$. We find using $J = 1/p$ that n is related to p through the relation

$$n = \frac{p}{p-2} \quad (26)$$

For $p = 2$ and $n = \infty$ the system is Z_2 equivalent to a fermionic gas. Also for $p = 3$ and 4 results $n = 3$ and 2, respectively, the system can be represented by fermions of fractional fermion number. On the other hand as $p \rightarrow \infty$ we have $n = 1$ system represents plane rotor model (XY model). This analysis suggests that we can go gradually from the Ising model (Z_2) to the XY model through changing p values.

As a fermion moving with angular momentum $J = 1/2$ departs from its pure fermionic character to Z_p spin system, it acquires the angular momentum $J = 1/p$. This leads to the change of angular momentum by

$$\Delta J = 1/2 - 1/p = \frac{p-2}{2p} = \frac{1}{2n} \quad (27)$$

which further indicate a shift in statistics

$$\Delta\theta = 2\pi\Delta J \quad (28)$$

If the composite fermion is in Z_2 spin system, changes to fermion in the ground state, there exists a shift in angular momentum by

$$\Delta J = 1/2 - 1/2 = 0$$

leading to change of statistics $\Delta\theta = 0$. Similarly for the composite fermion in the XY model, the deviation of angular momentum becomes

$$\Delta J = 1/2 - 1/\infty = 1/2$$

resulting the corresponding change in statistics of the Hall particles in the XY model to fermionic model is $\Delta\theta = \pi$.

These deviation in angular momentum can be visualized through the Shift quantum number in the frame work of Z_p spin system. Earlier we have studied [13] the Hall particles in hierarchies by attachment of extra strength of vortex and arrived at the Jain classification scheme from the equivalent relationship between fractional statistics and Z_p spin system where the composite fermion filling factor is given by

$$\nu_{CF}^{-1} = \theta/\pi + \tilde{m} = \frac{2}{p} + \tilde{m} = \frac{2mn+1}{n} \cong \frac{2mn' \pm 1}{n'} = 2\mu_{eff}/n' \quad (29)$$

where $\tilde{m}$ is an even number in connection with the flux attached with the composite particles.

In the light of Z_P spin system, the *Shift* in hierarchies from eqn.21 becomes

$$\mathcal{S} = n'(\frac{2}{p} + \tilde{m}) \mp (n' \pm 1) \quad (30)$$

in which using eqn. 27, for the release of the quanta the *Shift* is

$$\mathcal{S} = n'(\frac{2}{p} + \tilde{m}) - (n' \pm 1) = 2n'\Delta J + 2mn' \pm 1 = 2n'\Delta J + n'\nu_{CF}^{-1} = 2n'\Delta J + J_{CF} \quad (31)$$

On the other hand during acceptance of quanta, it becomes

$$\mathcal{S} = n'(\frac{2}{p} + \tilde{m}) + (n' \pm 1) = 2n'(\Delta J + 1) + 2mn' \pm 1 = 2n'(\Delta J + 1) + J_{CF} \quad (32)$$

It seems from above, that *Shift* in both the cases depend on angular momentum of composite fermion and it's change. The *Shift* during acceptance of quanta is more than that of release, by the term $2n'$ which yields no extra effect in the appearance of phase as seen from eqn.17.

A non-vanishing effect of *Shift* in 2D is obtained for its origin lying in the deviation of angular momentum of the Hall particles from the higher to lower Landau level. We further realize that the difference in angular momentum or the change in statistics is the very source of the effective flux N_ϕ (eqn5) present in any Landau level by either

$$2n'\Delta J \quad \text{or} \quad 2n'(\Delta J + 1) \quad (33)$$

depending on the respective accept or release of quanta by the Hall particles.

If the Hall fluid represent a plane rotor model(XY model), for $p \rightarrow \infty$ in the Z_p system the $\mathcal{S}$ becomes from eqn.(30)

$$\mathcal{S} = (2mn' \pm 1) \mp n' = n'\nu_{CF}^{-1} \mp n' = J_{CF} \mp n' \quad (34)$$

which indicates $\Delta J = 1/2$ and $\Delta\theta = \pi$.

On the other hand for Hall system behaving an Ising model,the *Shift* results

$$\mathcal{S} = n'\nu_{CF}^{-1} + n' \mp n' \quad (35)$$

following proper physics for the change in angular momentum $\Delta J = 0$, $\Delta\theta = \pi$ For acceptance or release of quanta the respective values of $\mathcal{S}$ are

$$\mathcal{S} = 2mn' \pm 1 = n'\nu_{CF}^{-1} = J_{CF} \quad \text{and} \quad \mathcal{S} = n'\nu_{CF}^{-1} + 2n' = J_{CF} + 2n' \quad (36)$$

imply identical effect since $2n'$ produces no extra effect in the topological phase. The effective flux associated with the respective XY model and Ising model becomes

$$n', 3n' \quad \text{and} \quad 0, 2n'$$

indicating a successive occurrence with the change of p values in Hall system.

We conclude by mentioning that we have given here a new definition of *Shift* from the point of view of angular momentum. It indicates the connection between fermionic and composite fermionic behavior of Hall particles in FQHE. A non-vanishing effect of *Shift* in 2D is shown here through the consideration of Hall particles as Z_p spin system. We have also shed light on the effective flux associated

with each Landau level. In addition we are successful in studying the physics of Hall system behaving as the Ising model and XY model through *Shift* quantum number and effective flux which re-open a new idea of FQHE from the topological aspect.

Acknowledgement

I like to express my gratitude specially to Prof. P. Bandyopadhyay and all the authors cited in my references.

References

- 1 K.Yu.Bliokh and Yu.P.Bilokh, quant-ph/0404144.
- 2 M.V.Berry; Proc.Roy.Soc.London; **A392**, 45 (1984), Proceedings of a NATO Advanced Research Workshop on Fundamental Aspects of Quantum Theory, held in Italy, Sep. 1985;
- 3 F.Wilczek; *Fractional Statistics, Anyon Superconductivity*, World Scientific, 1990.
- 4 P.Bandyopadhyay; *Geometry, Topology and Quantization* (kluwer, Dordrecht, The Netherlands, 1996).
- 5 D.Banerjee and P.Bandyopadhyay; Mod.Phys.Lett-**B8**, (1994), 1643.
- 6 D.Banerjee; Fort.Der.Physik **44** (1996) 323-370.
- 7 H.Kuratsuji and S.Iida; Phys.Rev **D37**, (1988), 441.
- 8 P.Bandyopadhyay; Int.J.Mod.Phys-**A15**, (2000), 1415.
- 9 F.D.M.Haldane, Phys.Rev.Lett **51**, 605 (1983).
- 10 D.Banerjee; Phys.Rev.**B58**, (1998), 4666.
- 11 X.G.Wen; Adv.in Phys. **41**, (1995) 405.
- 12 G. Goswami and P.Bandyopadhyay; J.Math.Phys. **33** (1992), 1090.
- 13 D.Banerjee and P.Bandyopadhyay; Phys. Lett-**A246**, (1998) 181.